

Entanglement and the $SU(2)$ phase states in atomic systems

M. Ali Can, Alexander A. Klyachko, and Alexander S. Shumovsky
Faculty of Science, Bilkent University, Bilkent, Ankara, 06533, Turkey
quant-ph/0202041, February, 2002

We show that a system of $2n$ identical two-level atoms interacting with n cavity photons manifests entanglement and that the set of entangled states coincides with the so-called $SU(2)$ phase states. In particular, violation of classical realism in terms of the GHZ and GHSH conditions is proved. We discuss a new property of entanglement expressed in terms of local measurements. We also show that generation of entangled states in the atom-photon systems under consideration strongly depends on the choice of initial conditions and that the parasitic influence of cavity detuning can be compensated through the use of Kerr medium.

I. INTRODUCTION

It has been recognized that entanglement phenomenon touches on the conceptual problems of reality and locality in quantum physics as well as the more technological aspects of quantum communications, cryptography, and computing. In particular, the methods of quantum key distribution in communication channels secured from eavesdropping are based on the use of entangled states [1,2,3,4,5] (for recent review, see Refs. [6,7]). In turn, the realization of quantum computer [8] is dependent on the ability to form entangled states of initially uncorrelated single-particle states [9].

In recent years, many successful experiments have been performed to verify the violation of Bell's inequalities and Greenberger-Horne-Zeilinger (GHZ) equality [10,11] and to develop the methods of engineered entanglement for quantum cryptography and quantum key distribution. In particular, the recent advances in the field of cavity QED and techniques of atom manipulation, trapping, and cooling enable a number of experiments which investigate the entanglement in the atomic systems (see [11,12,13,14,15,16,17] and references therein).

It has been shown recently [18] that a pure entangled state of two atoms can be obtained in an optical resonator through the exchange by a single photon. The main idea in Ref. [18] is that a single excitation of the system is either carried by a photon or shared between the atoms. If the photon can leak out from the resonator, the absence of photon counts in the process of continuous monitoring of the cavity decay can be associated with the presence of the pure entangled atomic state. The importance of this scheme is caused by the fact that its realization seems to be easily available with present experimental technique.

The main objective of this paper is to show that the entangled states in the "atoms plus photons" systems of the type discussed in [18] can be represented by the so-called $SU(2)$ phase states corresponding to the $SU(2)$ algebra of the odd "spin"

$$j = \frac{1}{2} \left[\binom{2n}{n} - 1 \right], \quad (1)$$

where $2n$ is the even number of atoms and $n = 1, 2, \dots$ is the number of cavity photons. In particular, the system considered in Ref. [18] corresponds to the phase states of "spin" $j = 1/2$. The $SU(2)$ phase states were introduced in [19] for an arbitrary spin and then generalized in [20,21] to the case of the $SU(2)$ subalgebra in the Weyl-Heisenberg algebra of photon operators (for recent review, see [22]). From the mathematical point of view, this is the system of

$$N = 2j + 1$$

qubits defined in the Hilbert space

$$\mathcal{H}_N = (\mathbf{C}^2)^{\otimes N}$$

with the componentwise action of $SU(2)^N$. In particular, we show that these states violate the classical realism and discuss their realization.

On the other hand, we will discuss a new condition of entanglement has been proposed recently [23]. Let us note in this connection that, in the usual treatment of entanglement, the entangled states of a two-component (in general, multi-component) system are considered as the nonseparable states with respect to the subsystems (e.g., see [24]). For example, if the individual components of a two-component system are described by the states $|\xi_i\rangle$ and $|\chi_i\rangle$, respectively, the state

$$|\psi_{ent}\rangle = \sum_i b_i |\xi_i\rangle \otimes |\chi_i\rangle, \\ \langle \xi_i | \xi_k \rangle = \langle \chi_i | \chi_k \rangle = \delta_{ik}, \quad \sum_i |b_i|^2 = 1$$

is entangled if $b_i \neq 0$ for at least two distinct values of the subscript i . From the mathematical point of view, the entanglement is caused by the combination of the superposition principle in quantum mechanics with the tensor product structure of the space of state of the two- or multi-component system [25].

Very often, the existence of entanglement is verified in terms of violation of Bell's inequalities and their generalizations [26,27,28,29,30,31]. Another way is based on the use of GHZ theorem [10]. A possibility to introduce more general inequalities is also discussed [32].

It should be noted that the use of Bell's inequalities and their numerous generalizations demonstrate nothing but the nonexistence of hidden variables. Moreover, it is possible to say that the unique, general, and mathematically correct definition of entanglement still does not exist (e.g., see Ref. [32]).

An interesting approach has been proposed recently [32]. Considering the state shared between Alice and Bob as a quantum communication channel, the authors of Ref. [32] concluded that the information in the case of entanglement is carried mostly by the correlations between the ends of the channel. These correlations manifest themselves by means of the local measurements on the sides of the channel [23].

Following [23], consider a composite system defined in the Hilbert space

$$\mathcal{H} = \bigotimes_{\ell} \mathcal{H}^{(\ell)}, \quad \ell \geq 2.$$

Let G be the group of dynamical symmetry of a subsystem in the composite system. Then the Hermitian operators g associated with representation of G in $\mathcal{H}^{(\ell)}$ define the set of local measurement on the corresponding side of the channel provided by a state $|\psi\rangle \in \mathcal{H}$. For example, in the case of $\mathcal{H}^{(\ell)} = \mathbf{C}^2$, corresponding to the EPR spin- $\frac{1}{2}$ system, $G = SU(2)$ and the set of local measurements can be specified by the infinitesimal generators of the $SL(2)$ group

$$\{g\} = \{\sigma_k^{(\ell)}\}, \quad k = 1, 2, 3,$$

which is the complexification of the $SU(2)$ group.

It was shown in [23] that the maximum correlation between the ends of the channel corresponds to the states such that

$$\forall g, \quad \langle g \rangle = 0.$$

This statement can be illustrated by the atoms-plus-photons systems under consideration. Consider first the set of two identical two-level atoms. Let $|e_\ell\rangle$ and $|g_\ell\rangle$ denote the excited and ground atomic states of the ℓ^{th} atom, respectively. Then, the entangled, maximum excited atomic states in the system "2 atoms plus 1 photon" considered in [18] are

$$|\psi_\pm\rangle = \frac{1}{\sqrt{2}}(|e_1g_2\rangle \pm |g_1e_2\rangle). \quad (2)$$

Then, the local measurement g can be described by the Pauli matrices

$$\begin{aligned} \sigma_1^{(\ell)} &= |e_\ell\rangle\langle g_\ell| + |g_\ell\rangle\langle e_\ell|, \\ \sigma_2^{(\ell)} &= -i|e_\ell\rangle\langle g_\ell| + i|g_\ell\rangle\langle e_\ell|, \\ \sigma_3^{(\ell)} &= |e_\ell\rangle\langle e_\ell| - |g_\ell\rangle\langle g_\ell|, \end{aligned} \quad (3)$$

i.e., by the infinitesimal generators of the algebra $SL(2)$. It is now a straightforward matter to check that

$$\forall i, \ell \quad \langle \psi_\pm | \sigma_i^{(\ell)} | \psi_\pm \rangle = 0, \quad (4)$$

where averaging is taken over the states (4). Another example is provided by the GHZ states [10]

$$|\psi_\pm^{(GHZ)}\rangle = \frac{1}{\sqrt{2}}(|e_1e_2e_3\rangle \pm |g_1g_2g_3\rangle), \quad (5)$$

corresponding to the maximum atomic excitation in the $3 + 3$ -system. It is easily seen that the averaging of the local operators (3) over (5) gives the same result as (4).

This property (4) can be used to define the entangled states.

We will show that the $SU(2)$ phase states of spin j defined by Eq. (1) in a $2n + n$ -type atom-photon system obey the nonseparability conditions, have the property (4), and manifest the violation of classical realism expressed in terms of the GHZ [10] and CHSH (Clauser-Horne-Shimony-Holt) [33] conditions.

The paper is organized as follows. In Sec. II, we consider the representation of the $SU(2)$ phase states. As a particular example, we examine the system of two identical two-level atoms, interacting with a single cavity photon and show that the maximum entangled atomic states of the Ref. [18] belong to the class of the $SU(2)$ phase states of spin $j = 1/2$. Let us stress that hereafter the maximum entanglement is defined in the usual way by the maximum of reduced entropy (e.g., see Refs. [23,25,27,32]). Then, we generalize this result on the case

of $2n + n$ system. As a nontrivial example we consider in Sec. III the system of four identical two-level atoms interacting with the two cavity photons. In this case, the set of entangled, maximum excited atomic states is provided by the six orthogonal $SU(2)$ phase states of spin $j = 5/2$. For these states, we prove violation of classical realism through the use of GHZ and CHSH conditions. In Sec. IV, we discuss how the entangled atomic states can be achieved in the process of steady-state evolution. In particular, we show that the maximum entanglement can be achieved if the initial state of the system contains the photons and does not contain the atomic excitations. We also show that the presence of the cavity detuning hampers the creation of pure entangled states and that the parasitic influence of detuning can be compensated through the use of the Kerr medium inside the cavity. Finally, in Sec. V, we briefly discuss the obtained results.

II. REPRESENTATION OF THE $SU(2)$ PHASE STATES

An arbitrary spin j can be described by the generators J_+, J_-, J_z of the $SU(2)$ algebra such that

$$\begin{aligned} [J_+, J_-] &= 2J_z, \quad [J_z, J_\pm] = \pm J_\pm, \\ J^2 &= J_z^2 + \frac{1}{2}(J_+J_- + J_-J_+) = j(j+1) \times \mathbf{1}, \end{aligned} \quad (6)$$

where $\mathbf{1}$ is the unit operator in the $2j + 1$ dimensional Hilbert space. Since

$$J_\pm = J_x \pm iJ_y,$$

it is possible to say that the generators J_+, J_-, J_z in (6) correspond to the Cartesian representation of the $SU(2)$ algebra. Following [19], one can introduce the representation in spherical coordinates via the polar decomposition of (6) of the form

$$J_+ = J_r \epsilon, \quad J_r = J_r^+, \quad \epsilon \epsilon^\dagger = \mathbf{1}, \quad (7)$$

where the Hermitian operator J_r corresponds to the radial contribution, while ϵ gives the exponential of the azimuthal phase operator. It is a straightforward matter to show that ϵ can be represented by the following $(2j + 1) \times (2j + 1)$ matrix

$$\epsilon = \begin{pmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 1 \\ e^{i\psi} & 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \quad (8)$$

in the $2j + 1$ -dimensional Hilbert space. Here ψ is an arbitrary real parameter (reference phase). The eigenstates of the operator (8)

$$\epsilon |\phi_n^{(j)}\rangle = e^{i\phi_n^{(j)}} |\phi_n^{(j)}\rangle, \quad n = 1, \dots, (2j + 1), \quad (9)$$

form the basis of the so-called phase states

$$|\psi_n^{(j)}\rangle = \frac{1}{\sqrt{2j+1}} \sum_{k=0}^{2j} e^{ik\phi_n^{(j)}} |\psi_k\rangle \quad (10)$$

dual with respect to the basis of individual states $|\psi_k\rangle$ of the Hilbert space.

As a physical example of some considerable interest, consider now the system of the two identical two-level atom interacting with the single cavity photon (see [18]). If the cavity photon is absorbed by either atom, the atomic subsystem can be observed in the following states

$$|\psi_1\rangle = |e_1g_2\rangle, \quad |\psi_2\rangle = |g_1e_2\rangle, \quad (11)$$

where $|e_1g_2\rangle = |e_1\rangle \otimes |g_2\rangle$ and $|e\rangle$ and $|g\rangle$ denote the excited and ground atomic states, respectively. The subscript marks the atom. Using the atomic basis (11), we can construct the following representation of the $SU(2)$ algebra:

$$\begin{aligned} J_+ &= |e_1g_2\rangle\langle g_1e_2|, \quad J_- = |g_1e_2\rangle\langle e_1g_2|, \\ J_3 &= \frac{1}{2}(|e_1g_2\rangle\langle e_1g_2| - |g_1e_2\rangle\langle g_1e_2|). \end{aligned} \quad (12)$$

This representation formally corresponds to (6) at the spin $j = 1/2$. Then, the corresponding exponential of the phase operator (8) takes the form

$$\epsilon = |e_1g_2\rangle\langle g_1e_2| + e^{i\psi}|g_1e_2\rangle\langle e_1g_2|. \quad (13)$$

In turn, the phase states (9) and (10) are

$$\begin{aligned} |\phi_{\pm}\rangle &= \frac{1}{\sqrt{2}}(|e_1g_2\rangle + e^{i\phi_{\pm}}|g_1e_2\rangle), \\ \phi_{\pm} &= \psi/2 + (1 \mp 1)\pi/2. \end{aligned} \quad (14)$$

It is easily seen that the phase states (14) form the set of entangled atomic states in the two-atom system under consideration. Definitely, these states obey the nonseparability condition. It is also seen that (14) coincides with the maximally entangled states (2) of Ref. [18] when the reference phase $\psi = 0$.

Consider now a general $2n+n$ system at $n \geq 1$. Then, the maximum excited atomic states

$$|\psi_i\rangle = |\{e\}_n, \{g\}_n\rangle, \quad (15)$$

can be used to construct a representation of the $SU(2)$ algebra (6) of spin j defined in (1). Here $i = 1, 2, \dots, N$ and

$$N = 2j + 1 = \binom{2n}{n}$$

is the total number of such a states. In the basis (15), we can construct the polar decomposition of the $SU(2)$ algebra of spin (1) and the corresponding exponential of the phase operator (8) and the phase states (10). Let us rename the states (15) as follows

$$|\psi_k\rangle \rightarrow |\psi_{k'}\rangle, \quad k' \equiv k - 1 = 0, \dots, N - 1.$$

Then, the $SU(2)$ phase states (10) take the form

$$|\phi_k\rangle = \frac{1}{\sqrt{N}} \sum_{k'=0}^{N-1} e^{ik'\phi_k} |\psi_{k'}\rangle, \quad (16)$$

where

$$\phi_k = (\psi + 2k\pi)/N.$$

These states (16) form a basis dual with respect to (15) and spanning the Hilbert space of the maximum excited atomic states in the $2n+n$ system under consideration. By construction, the phase states (16) are nonseparable with respect to contributions of individual atoms and thus entangled [24]. Let us stress that the choice of the phase factors in (16) is irrelevant to entanglement, which holds for arbitrary phase factors. This choice is caused by the aspiration for getting the dual with respect to (15) basis of entangled states.

It is easily seen that the states (16) obey the condition (4). In fact, the action of the flip-operators $\sigma_{1,2}^{(\ell)}$ in (3) on the states (16) leads to the change of the number of either excited or de-excited atoms:

$$\sigma_{1,2}^{(\ell)} |\psi_k\rangle \rightarrow \begin{cases} |\{e\}_{n-1}, \{g\}_{n+1}\rangle & \ell \in \{g\} \\ |\{e\}_{n+1}, \{g\}_{n-1}\rangle & \ell \in \{e\} \end{cases}$$

and therefore $\langle \sigma_{1,2}^{(\ell)} \rangle = 0$ in the case of averaging over the states (16). Since each state (15) contains equal number of excited and de-excited atoms, the action of the parity operator in (3) on the phase states (16) should lead to the state which differ from (16) by the multiplication of a certain n terms by the factor of -1 . Hence

$$\langle \sigma_3^{(\ell)} \rangle = \frac{1}{N} \left(\sum_{i=1}^{N/2} 1 - \sum_{i=N/2+1}^N 1 \right) = 0.$$

By construction, N is always an even number.

Thus, the $SU(2)$ phase states (16), corresponding to the maximum excited atomic states in the $2n+n$ system, are entangled because they are nonseparable and, at the same time, obey the condition (4) for the local measurements. In the next Section, we show that the states (16) manifest violation of classical realism as well.

Before we begin to discuss this subject, let us note that the $SU(2)$ phase states of the atomic system under consideration with integer spin do not provide the entanglement. Consider as an example the system of three identical two-level atoms, interacting with a single cavity photon. There are the three excited atomic states

$$|e_1g_2g_3\rangle, \quad |g_1e_2g_3\rangle, \quad |g_1g_2e_3\rangle \quad (17)$$

and the three dual phase states of the type of (16)

$$|\psi_k\rangle = \frac{1}{\sqrt{3}}(|e_1g_2g_3\rangle + e^{i\phi_k}|g_1e_2g_3\rangle + e^{2i\phi_k}|g_1g_2e_3\rangle). \quad (18)$$

It is clear that the states (18) are the phase states of spin $j = 1$. Here

$$\phi_k = (\psi + 2k\pi)/3, \quad k = 0, 1, 2.$$

It is easily seen that the phase states (18) cannot be factorized with respect to atoms. At the same time, the average of the parity operator $\sigma_3^{(\ell)}$ in (3) over the states (18) is

$$\forall k, \ell \quad \langle \psi_k | \sigma_3^{(\ell)} | \psi_k \rangle = -\frac{1}{3},$$

although the averages of the flip-operators are

$$\forall k, \ell \quad \langle \psi_k | \sigma_{1,2}^{(\ell)} | \psi_k \rangle = 0.$$

Thus, the nonseparable states (18) do not obey the condition (4). At the same time, these states do not manifest the maximum entanglement as well. Let us stress that the nonseparability is not a sufficient condition of maximum entanglement [24]. For example, from the measurement of the state of the first atom we can only learn that either the atoms 2 and 3 are both in the ground state with reliability or they are in the two-atom entangled state of the type discussed in [18]. Similar result can be obtained for the system of three atoms, interacting with two cavity photons. The only maximum entangled state of the system of three atoms is provided by the superposition of GHZ states (5).

III. THE 4 + 2-SYSTEM

To show that the phase states (16) of a $2n + n$ system violate the classical realism, consider the system of four identical two-level atoms interacting with two cavity photons. The maximum excited atomic states at $n = 2$ are

$$\begin{aligned} &|e_1 e_2 g_3 g_4\rangle, \quad |e_1 g_2 e_3 g_4\rangle, \quad |e_1 g_2 g_3 e_4\rangle, \\ &|g_1 e_2 e_3 g_4\rangle, \quad |g_1 e_2 g_3 e_4\rangle, \quad |g_1 g_2 e_3 e_4\rangle. \end{aligned} \quad (19)$$

These orthonormal states form the six-dimensional basis of the Hilbert space in which the representation of the generators (6) has the form

$$\begin{aligned} J_+ &= \sqrt{5}|e_1 e_2 g_3 g_4\rangle\langle e_1 g_2 e_3 g_4| + \sqrt{8}|e_1 g_2 e_3 g_4\rangle\langle e_1 g_2 g_3 e_4| \\ &\quad + 3|e_1 g_2 g_3 e_4\rangle\langle g_1 e_2 e_3 g_4| + \sqrt{8}|g_1 e_2 e_3 g_4\rangle\langle g_1 e_2 g_3 e_4| \\ &\quad + \sqrt{5}|g_1 e_2 g_3 e_4\rangle\langle g_1 g_2 e_3 e_4|, \\ J_3 &= \frac{5}{2}|e_1 e_2 g_3 g_4\rangle\langle e_1 e_2 g_3 g_4| + \frac{3}{2}|e_1 g_2 e_3 g_4\rangle\langle e_1 g_2 e_3 g_4| \\ &\quad + \frac{1}{2}|e_1 g_2 g_3 e_4\rangle\langle e_1 g_2 g_3 e_4| - \frac{1}{2}|g_1 e_2 e_3 g_4\rangle\langle g_1 e_2 e_3 g_4| \\ &\quad - \frac{3}{2}|g_1 e_2 g_3 e_4\rangle\langle g_1 e_2 g_3 e_4| - \frac{5}{2}|g_1 g_2 e_3 e_4\rangle\langle g_1 g_2 e_3 e_4| \end{aligned}$$

By construction, they describe the spin $j = 5/2$ system. In turn, the exponential of the phase operator (8) takes the form

$$\begin{aligned} \epsilon &= |e_1 e_2 g_3 g_4\rangle\langle e_1 g_2 e_3 g_4| + |e_1 g_2 e_3 g_4\rangle\langle e_1 g_2 g_3 e_4| \\ &\quad + |e_1 g_2 g_3 e_4\rangle\langle g_1 e_2 e_3 g_4| + |g_1 e_2 e_3 g_4\rangle\langle g_1 e_2 g_3 e_4| \\ &\quad + |g_1 e_2 g_3 e_4\rangle\langle g_1 g_2 e_3 e_4| + e^{i\psi}|g_1 g_2 e_3 e_4\rangle\langle e_1 e_2 g_3 g_4|. \end{aligned}$$

Then, the six phase states (9) have the form (16) with $N = 6$ and

$$\phi_k = \frac{\psi}{6} + \frac{k\pi}{3}, \quad k = 0, 1, \dots, 5. \quad (20)$$

As well as (16), these states are nonseparable and hence entangled and obey the condition (4) for local variables.

To show that these phase states violate the classical realism, let us first represent the states (16) at $N = 6$ in the following way

$$|\phi_k\rangle = \frac{1}{\sqrt{3}}(|\chi_{1k}\rangle + e^{i\phi_k}|\chi_{2k}\rangle + e^{2i\phi_k}|\chi_{3k}\rangle), \quad (21)$$

where

$$\begin{aligned} |\chi_{1k}\rangle &= \frac{1}{\sqrt{2}}(|e_1 e_2 g_3 g_4\rangle + e^{5i\phi_k}|g_1 g_2 e_3 e_4\rangle), \\ |\chi_{2k}\rangle &= \frac{1}{\sqrt{2}}(|g_1 e_2 e_3 g_4\rangle + e^{3i\phi_k}|e_1 g_2 g_3 e_4\rangle), \\ |\chi_{3k}\rangle &= \frac{1}{\sqrt{2}}(|g_1 e_2 g_3 e_4\rangle + e^{i\phi_k}|e_1 g_2 e_3 g_4\rangle). \end{aligned} \quad (22)$$

It is easily seen that each set of six states $|\chi_{pk}\rangle$ with $p = 1, 2, 3$ and $k = 0, \dots, 5$ consists of the nonseparable and hence entangled states. Consider, for example, the states $|\chi_{1k}\rangle$ in (22). Because of the definition of the phase angle ϕ_k at $N = 6$, they consist of the three sets of the pairwise orthogonal states

$$\{|\chi_{10}\rangle, |\chi_{13}\rangle\}, \quad \{|\chi_{11}\rangle, |\chi_{14}\rangle\}, \quad \{|\chi_{12}\rangle, |\chi_{15}\rangle\}.$$

It is also seen that the second and third sets here are obtained from the first set by the successive rotations of the reference frame.

Now the violation of classical realism can be proved through the use of the GHZ theorem [10]. Consider first the state $|\chi_{10}\rangle$ in (22). It is easy to verify that this state obey the conditions

$$\forall i, \ell \quad \bigotimes_{\ell=1}^4 \sigma_i^{(\ell)} |\chi_{10}\rangle = |\chi_{10}\rangle \quad (23)$$

and

$$\begin{aligned} \sigma_1^{(1)} \sigma_1^{(2)} \sigma_2^{(3)} \sigma_2^{(4)} |\chi_{10}\rangle &= -|\chi_{10}\rangle, \\ \sigma_2^{(1)} \sigma_2^{(2)} \sigma_1^{(3)} \sigma_1^{(4)} |\chi_{10}\rangle &= -|\chi_{10}\rangle, \\ \sigma_1^{(1)} \sigma_2^{(2)} \sigma_1^{(3)} \sigma_2^{(4)} |\chi_{10}\rangle &= |\chi_{10}\rangle, \\ \sigma_1^{(1)} \sigma_2^{(2)} \sigma_2^{(3)} \sigma_1^{(4)} |\chi_{10}\rangle &= |\chi_{10}\rangle, \\ \sigma_2^{(1)} \sigma_1^{(2)} \sigma_2^{(3)} \sigma_1^{(4)} |\chi_{10}\rangle &= |\chi_{10}\rangle, \\ \sigma_2^{(1)} \sigma_1^{(2)} \sigma_1^{(3)} \sigma_2^{(4)} |\chi_{10}\rangle &= |\chi_{10}\rangle. \end{aligned} \quad (24)$$

It is possible to say that these equalities (23) and (24) express a kind of EPR "action at distance" in the maximum excited states of the system of four atoms interacting with two photons. In other words, the correlations represented by (23) and (24) permit us determine in a unique way the state of the fourth atom via measurement of the states of other three atoms.

The operator equalities (23) and (24) can be used to obtain the relations similar to those in the GHZ theorem. Following [10], we have to assign the classical quantities $m_i^{(\ell)}$ to the local operators. Here

$$m_1^{(\ell)}, m_2^{(\ell)} = \pm 1.$$

Then, it follows from (23) that

$$\prod_{\ell=1}^4 m_1^{(\ell)} = 1. \quad (25)$$

At the same time, it follows from (24) that

$$[\sigma_1^{(1)} \sigma_1^{(2)} \sigma_2^{(3)} \sigma_2^{(4)}][\sigma_1^{(1)} \sigma_2^{(2)} \sigma_1^{(3)} \sigma_2^{(4)}][\sigma_1^{(1)} \sigma_2^{(2)} \sigma_2^{(3)} \sigma_1^{(4)}]|\chi_{10}\rangle = -|\chi_{10}\rangle.$$

Employing the classical variables instead of the local operators allows this to be cast into the form

$$(m_1^{(1)})^3 m_1^{(2)} (m_2^{(2)})^2 m_1^{(3)} (m_2^{(3)})^2 m_1^{(4)} (m_2^{(4)})^2 = -1.$$

Since $(m_1^{(\ell)})^2 = (m_2^{(\ell)})^2 = 1$, we get an equivalent equality

$$m_1^{(1)} m_1^{(2)} m_1^{(3)} m_1^{(4)} = -1,$$

which contradicts (25). Hence, the state $|\chi_{10}\rangle$ in (22) obey the GHZ theorem. Similar result can be obtained for all other states in (22) and hence, for the phase states (21).

Our consideration so far have applied to the local measurements touching on a single atom. We now note that the phase states (21) allow another kind of entanglement in the case of pairwise measurement. Consider again the state $|\chi_{10}\rangle$ in (22) and assume that the measurements a and b corresponds to a pair of atoms:

$$\begin{aligned} a &= \cos \theta_a |e_1 e_2\rangle \langle e_1 e_2| + \sin \theta_a (|e_1 e_2\rangle \langle g_1 g_2| \\ &\quad + |g_1 g_2\rangle \langle e_1 e_2| - \cos \theta_a |g_1 g_2\rangle \langle g_1 g_2|, \\ b &= \cos \theta_b |e_3 e_4\rangle \langle e_3 e_4| + \sin \theta_b (|e_3 e_4\rangle \langle g_3 g_4| \\ &\quad + |g_3 g_4\rangle \langle e_3 e_4| - \cos \theta_b |g_3 g_4\rangle \langle g_3 g_4|. \end{aligned} \quad (26)$$

Assume now that we make the two measurements a and a' with the angles $\theta_1 = \pi$ and $\theta'_a = \pi/2$ and the two more measurements b and b' with the angles $\theta'_b = -\theta_b$, respectively. Then, the averaging over the state $|\chi_{10}\rangle$ gives

$$\langle ab \rangle = \langle ab' \rangle = \cos \theta_b, \quad \langle a'b \rangle = \sin \theta_b = -\langle a'b' \rangle.$$

Employing the CHSH inequality [33]

$$|\langle ab \rangle + \langle a'b \rangle + \langle a'b' \rangle - \langle ab' \rangle| \leq 2 \quad (27)$$

then gives

$$|\cos \theta_b - \sin \theta_b| \leq 1.$$

Violation of this inequality and hence, of the classical realism occurs at small negative θ_b , when we can put

$$|\cos \theta_b - \sin \theta_b| \sim 1 + |\theta_b| > 1.$$

Similar consideration can be done for all over states in (22) through the use of proper pairwise measurements. At the same time, the phase states (21) do not manifest entanglement with respect to the pairwise measurements.

The phase states (16) for the $6+3, 8+4, \dots$ systems, corresponding to the spin (1) equal to $19/2, 69/2, \dots$, respectively, can be considered as above.

IV. INITIAL CONDITIONS AND ATOMIC ENTANGLEMENT

It is clear that the evolution of the $2n+n$ system strongly depends on the choice of initial conditions. To trace the proper choice leading to the atomic entanglement, let us ignore the relaxation processes. Then, the steady-state evolution of the $2n+n$ system under consideration is governed by the Hamiltonian

$$H = \Delta a^+ a + \omega_0 \mathcal{N} + \gamma \sum_{\ell} (R_{\ell}^+ a + a^+ R_{\ell}). \quad (28)$$

Here Δ is the cavity detuning, ω_0 is the atomic transition frequency, γ is the atom-field coupling constant, operators a and a^+ describe the cavity photons,

$$\mathcal{N} = a^+ a + \sum_{\ell} |e_{\ell}\rangle \langle e_{\ell}| \bigotimes_{\ell' \neq \ell} \mathbf{1}^{(\ell')},$$

and the atomic operators are defined as follows

$$R_{\ell}^+ = |e_{\ell}\rangle \langle g_{\ell}| \bigotimes_{\ell' \neq \ell} \mathbf{1}^{(\ell')}.$$

Here $\mathbf{1}^{(\ell)}$ denotes the unit operator in the two-dimensional Hilbert space of the ℓ^{th} atom. It is seen that $[\mathcal{N}, H] = 0$. It is also seen that the atomic operators are similar, in a certain sense, to the local operators (3). In fact

$$R_{\ell}^{\pm} = \frac{\sigma_1^{(\ell)} \pm i \sigma_2^{(\ell)}}{2}.$$

Consider first the case of two atoms and single cavity photon when $\ell = 1, 2$ and the Hamiltonian (28) coincides with that of Ref. [18]. For simplicity, we use here the

same coupling constant γ for both atoms. Our consideration can easily be generalized on the case of coupling constant depending on the atomic position. Let us note that, in the case of only two atoms, the Hamiltonian (28) can be represented as follows

$$H \rightarrow H_\phi = \Delta a^\dagger a + \omega_0 \mathcal{N}_\phi + \gamma \sqrt{2} (\mathcal{R}^\dagger a + a^\dagger \mathcal{R}), \quad (29)$$

where

$$\mathcal{N}_\phi = a^\dagger a + \sum_{k=\pm 1} |\phi_k\rangle \langle \phi_k|$$

and

$$\mathcal{R}^\dagger = |\phi_+\rangle \langle g_1 g_2|.$$

Here $|\phi_\pm\rangle$ denote the phase states (14).

Using the Hamiltonian (29) as the generator of evolution, for the time dependent wave function we get

$$\begin{aligned} |\Psi(t)\rangle &= e^{-iH_\phi t} |\Psi(0)\rangle = [C_-(t) |\phi_-\rangle \\ &+ C_+(t) |\phi_+\rangle] \otimes |0\rangle_{ph} + C(t) |g_1 g_2\rangle \otimes |1\rangle_{ph}, \end{aligned} \quad (30)$$

where $|\dots\rangle_{ph}$ denotes the states of the cavity field. The coefficients $C_\pm(t)$ and $C(t)$ in (30) are completely determined by the initial conditions and normalization condition.

It is easily seen that the state $|\phi_-\rangle \otimes |0\rangle_{ph}$ is the eigenstate of the Hamiltonian (29). Hence, at

$$C_-(0) = 1, \quad C_+(0) = C(0) = 0,$$

the atomic phase state $|\phi_-\rangle$ in (14) provides the stationary, maximum entangled atomic state in the system under consideration [18]. At the same time, it is not very clear how to prepare such a state.

Therefore we consider a more realistic initial state provided by excitation of either atom, while the cavity field is in the vacuum state. To realize such a state, we can assume, for example, that one of the atoms (initially de-excited) is trapped in the cavity, while the second atom (initially excited) slowly passes through the cavity like in the experiments discussed in [14,15]. assume for definiteness that

$$|\Psi(0)\rangle = |e_1 g_2\rangle \otimes |0\rangle_{ph}. \quad (31)$$

Then, the coefficients of the wave function (30) take the form

$$\begin{aligned} C_-(t) &= \frac{1}{\sqrt{2}} e^{-i\omega_0 t}, \\ C_+(t) &= \frac{1}{\sqrt{2}} \left(\cos \Omega t + \frac{i\Delta}{2\Omega} \sin \Omega t \right) e^{-i(\omega_0 + \Delta/2)t}, \\ C(t) &= -\frac{i\gamma}{\Omega} e^{-i(\omega_0 + \Delta/2)t} \sin \Omega t, \end{aligned}$$

where $\Omega = [2\gamma^2 + (\Delta/2)^2]^{1/2}$. At first site, the probabilities

$$P_\pm(t) = |\langle 0|_{ph} \otimes \langle \phi_{pm} | \Psi(t) \rangle|^2 = |C_\pm(t)|^2$$

to observe the states (14) corresponding to the maximum atomic entanglement, are

$$\begin{aligned} P_-(t) &= \frac{1}{2}, \\ P_+(t) &= \frac{\Delta^2}{8\Omega^2} + \frac{\gamma^2}{\Omega^2} \cos^2 \Omega t \leq \frac{1}{2}, \end{aligned}$$

respectively. At the same time, the absence of photon counts, which is considered in [18] as a sign of the atomic entanglement, corresponds here to the case when both probabilities $P_\pm(t_k) = 1/2$ at a certain time t_k . In other words, the mutually orthogonal entangled states (14) have the same probability to be observed at $t = t_k$. This means that there is no atomic entanglement at all but we definitely know which atom is in the excited state.

Consider one more realistic initial state when both atoms are trapped in the cavity in de-excited state, while the cavity field contains a photon:

$$|\Psi(0)\rangle = |g_1 g_2\rangle \otimes |1\rangle_{ph}. \quad (32)$$

Then, for all times we get $C_-(t) = 0$ and

$$\begin{aligned} C_+(t) &= -\frac{i\gamma\sqrt{2}}{\Omega} e^{-i(\omega_0 + \Delta/2)t} \sin \Omega t, \\ C(t) &= \left(\cos \Omega t - \frac{i\Delta}{2\Omega} \sin \Omega t \right) e^{-i(\omega_0 + \Delta/2)t}. \end{aligned}$$

Hence, under this initial condition, the entangled state $|\phi_-\rangle$ cannot be achieved at all, while the second entangled state $|\phi_+\rangle$ in (14) can be achieved. It is seen that, in the case of initial state (32), the probability to detect the photon is

$$P_{ph}(t) = |C(t)|^2 = \cos^2 \Omega t + \frac{\Delta^2}{4\Omega^2} \sin^2 \Omega t.$$

This expression takes the minimum value

$$\min P_{ph} = P_{ph}(t_m) = \frac{\Delta^2}{4\Omega^2}$$

at $t = t_m = \pi(2m+1)/2\Omega$, $m = 0, 1, \dots$. At the same time t_m , the probability to have the entangled atomic state $|\phi_+\rangle$ takes the maximum value

$$P_+(t_m) = |C_+(t_m)|^2 = \frac{2\gamma^2}{2\gamma^2 + (\Delta/2)^2}.$$

It is seen that the pure atomic entanglement with $P_+(t_m) = 1$ is realized at $t = t_m$ only in the absence of the cavity detuning when $\Delta \rightarrow 0$.

The parasitic influence of the cavity detuning can be compensated through the use of Kerr medium filling the cavity. In this case, the Hamiltonian (28) should be supplemented by the term

$$H_\kappa = \kappa(a^\dagger a)^2,$$

which leads to the following renormalization of the Rabi frequency

$$\Omega \rightarrow \Omega_\kappa = \sqrt{2\gamma^2 + (\Delta + \kappa)^2}/4.$$

Then, the proper choice of the Kerr parameter $\kappa = -\Delta$ should lead to the pure entangled atomic state $|\phi_+\rangle$ at a certain times.

Consider now the case of four atoms and two photons. In contrast to the previous case, neither phase state in (21) is an eigenstate of the Hamiltonian (28). Then, the choice of the initial state either as a state with two excited atoms or as a state with one excited atom plus cavity photon does not lead to a pure atomic entanglement. As in the case of two atoms, the pure atomic entanglement can be reached under the choice of the state with the absence of the atomic excitations in the initial state. The influence of the cavity detuning can be compensated by the presence of Kerr medium as well as in the case of two atoms.

V. CONCLUSION

Let us briefly discuss the obtained results. For the system of two identical two-level atoms interacting with a single photon has been proposed in [18] it is shown that the maximum entangled atomic states are represented by the $SU(2)$ phase states of spin $1/2$. Moreover, it is shown that the $SU(2)$ phase states of the half-integer spin j (1) form a certain class of maximum entangled atomic states in the system of $2n$ atoms interacting with n photons. In particular, the violation of classical realism is shown.

It should be noted in this connection that the above considered $SU(2)$ phase states do not represent a unique way to construct the maximum entangled states in the multi-atom systems and that some other symmetries, for example the $SU(N)$ can be considered as well. Moreover, in some cases the $SU(2)$ phase states cannot be used to determine the maximum entangled states at all. Consider for example the case of two identical two-level atoms interacting with two photons, when the atomic subsystem can be specified by the four states

$$|e_1e_2\rangle, \quad |e_1g_2\rangle, \quad |g_1e_2\rangle, \quad |g_1g_2\rangle.$$

By performing a similar analysis to that described in section II, it is easy to construct the corresponding set of the $SU(2)$ phase states

$$|\phi_k\rangle = \frac{1}{2}(|e_1e_2\rangle + e^{i\phi_k}|e_1g_2\rangle + e^{2i\phi_k}|g_1g_2\rangle + e^{3i\phi_k}|g_1e_2\rangle),$$

$$\phi_k = \frac{\psi}{4} + \frac{k\pi}{2}, \quad k = 0, 1, 2, 3,$$

which do not manifest the maximum entanglement. At the same time, the general criterion (4) permits us to determine infinitely many maximum entangled states in

this case [23]. An example is provided by the following set of orthonormal maximum entangled states

$$|\psi_1\rangle = \frac{1}{2}(|e_1e_2\rangle + |g_1g_2\rangle + i|e_1g_2\rangle + i|g_1e_2\rangle),$$

$$|\psi_2\rangle = \frac{1}{2}(|e_1e_2\rangle - |g_1g_2\rangle - i|e_1g_2\rangle + i|g_1e_2\rangle),$$

$$|\psi_3\rangle = \frac{1}{2}(i|e_1e_2\rangle + i|g_1g_2\rangle + |e_1g_2\rangle + |g_1e_2\rangle),$$

$$|\psi_4\rangle = \frac{1}{2}(-i|e_1e_2\rangle + i|g_1g_2\rangle + |e_1g_2\rangle - |g_1e_2\rangle).$$

In fact, the Eq. (4) gives a general condition [23], while the $SU(2)$ phase states can manifest the maximum entanglement only under a certain condition (special choice of the effective spin (1)).

Nevertheless, the $SU(2)$ phase states considered in sections II and III represent an important example of the atomic entangled states. First of all, they can be easily realized in the atomic systems in a cavity. In fact, these states have a simple physical meaning. In addition to (9), the $SU(2)$ phase states can be defined to be the eigenstates of the Hermitian "cosine" operator [19,22]

$$C = \frac{1}{2}(\epsilon + \epsilon^\dagger),$$

where ϵ is defined by Eq. (8). This operator C can be considered as a "Hamiltonian", describing the correlations between the different atoms. For example, in the case of the two atoms interacting with the single photon, the operator C takes the form

$$C = \sigma_+^{(1)}\sigma_-^{(2)} + \sigma_-^{(2)}\sigma_+^{(1)}, \quad (33)$$

where

$$\sigma_\pm^{(\ell)} = \frac{\sigma_1^{(\ell)} \pm i\sigma_2^{(\ell)}}{2}.$$

The operator structure of (33) coincides with that of the so-called model of plane rotator, which is a particular case of the Heisenberg model of ferromagnetism widely used in statistical physics [35] and in quantum information theory [36].

Let us also stress that the $SU(2)$ phase states similar to those considered in sections II and III, have been discussed recently in the context of quantum coding [37].

It is also known that the $SU(2)$ phase states have direct connection with the quantum description of polarization of spherical photons emitted by the multipole transitions in atoms and molecules [21,22,38]. Therefore, the polarization entanglement of photons can be examined in direct analogy to the above discussed atomic entanglement [39]. At the same time, the consideration of spherical photons requires the use of more quantum degrees of freedom. Consider as an example the cascade decay of a two-level atom specified by the transition [40]

$$|J = 2, m = 0\rangle \rightarrow |J' = 0, m' = 0\rangle.$$

Here J, J' and m, m' denote the angular momentum and projection of the angular momentum of the excited and ground atomic states, respectively. This transition gives rise to an entangled photon twins [40]. Each photon carries spin 1, but because of the conservation of the angular momentum in the process of radiation, the sum of projections of the angular momenta of the two photons should be equal to zero. Denoting the state of a photon with given m by $|m\rangle$, we get the three possible states of the photon subsystem

$$|+1\rangle \otimes |-1\rangle, \quad |0\rangle \otimes |0\rangle, \quad |-1\rangle \otimes |+1\rangle.$$

These three "individual" states can be used to construct the dual basis of the $SU(2)$ phase states [21]

$$\begin{aligned} |\phi_k\rangle &= \frac{1}{\sqrt{3}}(|+1\rangle \otimes |-1\rangle + e^{i\phi_k}|0\rangle \otimes |0\rangle \\ &\quad + e^{2i\phi_k}|-1\rangle \otimes |+1\rangle), \\ \phi_k &= \frac{\psi + 2k\pi}{3}, \quad k = 0, 1, 2 \end{aligned} \quad (34)$$

similar to (18). It can be easily seen that these states manifest the maximum entanglement.

Similar entangled states have been discussed in the context of the so-called biphoton excitations [41] (photon pairs in symmetric Fock states). They can also be used in quantum cryptography [42].

Let us stress that the general condition of the type of (4) is also valid in the case of states (34). However, the definition of local measurement should be changed in this case. Because of the number of degrees of freedom per photon is equal to three, the Hermitian operators associated with the $SU(3)$ group should be considered instead of the infinitesimal generators of the $SL(2)$ group. For example, the set of the Stokes operators of the Ref. [21], corresponding to the representation of the $SU(3)$ subalgebra in the Weyl-Heisenberg algebra of spherical photons, can be used to define the complete set of local measurements in this case.

It is shown in section IV that the realization of a pure atomic entanglement in the $2n + n$ -type atom + photon systems strongly depends on the choice of initial state. Viz, the entangled states can be reached in the process of steady-state evolution only if all $2n$ atoms are initially in the de-excited states, while the cavity contains just n photons. This condition has an intuitively clear explanation: the excitations of different atoms have the same probability and therefore each photon in the $2n + n$ -system is shared with a couple of atoms.

It is also shown in section IV that the presence of the cavity detuning hampers the creation of a pure entangled atomic state. This negative effect can be compensated through the use of Kerr medium in the cavity.

We now note that the practical realization of a long-lived, maximum entanglement in a quantum mechanical system strongly depends on the interaction between this system and environment. The point is that the state of a

closed quantum mechanical system changes periodically, providing the maximum entanglement as an instant event only at a certain times (see section IV). Such a periodicity is caused by a finite number of degrees of freedom in the system. To destruct such a periodicity, it is necessary to connect the system to a "heat bath", which would tune in the system to a required state. In Ref. [18], it has been proposed to support the atomic entanglement by the cavity losses. In this case, the absence of the photon counting outside the cavity can be associated with the existence of the entangled atomic state in the cavity.

Let us stress that an advantage of the use of the $SU(2)$ phase states as the maximum entangled atomic states consists in the simple preparation of the initial states discussed in section IV.

In view of realization of atomic entanglement with present experimental technique, it seems to be more convenient if the existence of entangled state in a cavity would manifest itself via a signal photon rather than the absence of photon leakage from the cavity. In this case, there should be at least the two modes such that one of them (the cavity mode) provides the correlation between the atoms, while the second can freely leave the resonator to signalize the existence of the entanglement. Such a process can be realized through the use of the Raman process in atoms shown in Fig. 1 (e.g., see [43]). Here the dipole transitions are allowed between the levels 1 and 2 and 2 and 3, while forbidden between 1 and 3 because of the parity conservation. In the simplest case, we should assume that the two identical atoms of this type are located in a cavity, which has a very high quality with respect to the pumping mode ω_P , while the Stokes photons with frequency ω_{Sk} can leak away freely.

Assume that the atoms are initially in the ground state 1, the Stokes field is in the vacuum state, and the pump field consists of a single photon. The evolution of the system can lead to the absorption of the cavity photon by either atom with further emission of the Stokes photon, which leaves the cavity. After that, the atoms are in entangled state, corresponding to the excitation of the atomic level 3 shared between the atoms. Since the inverse process cannot be realized without assistance of the Stokes photon, such a state represents a durable atomic entangled state.

It is clear that the above consideration of the atomic entanglement in the multi-atom system can be generalized with easy on the case of Raman process in atoms. In other words, the $SU(2)$ phase states similar to (16) form the class of the maximum entangled atomic states in the case of Raman-type processes in the three-level atoms as well. An evident advantage of the use of the Raman process is the long-lived maximum entanglement in atomic subsystem.

One of the authors (A.S.) would like to thank Dr. A. Beige, Prof. P.L. Knight, Prof. A. Vourdas, and Prof. A. Zeilinger for useful discussions.

-
- [1] C.H. Bennet and G. Brassard, in *Proceedings of IEEE International Conference on Computers, systems and Signal Processing, Bangalore, India* (IEEE New York, 1984).
- [2] A.K. Ekert, Phys. Rev. Lett. **68**, 661 (1991).
- [3] C.H. Bennet, G. Brassard, and A.K. Ekert, Sci. America (Int. Ed.) **276**, 50 (1992).
- [4] A.K. Ekert, J.G. Rarity, P.G. Tapster, and G.M. Palma, Phys. Rev. Lett. **69**, 1293 (1992).
- [5] C.H. Bennet, F. Bessette, G. Brassard, L. Salivan, and J. Smolin, J. Cryptography **5**, 3 (1992).
- [6] D. Bouwmeester, A.K. Ekert, and A. Zeilinger, editors, *The Physics of Quantum Information* (Springer-Verlag, Berlin, 2000).
- [7] P. Tombesi and O. Hirota, editors, *Quantum Communications, Computing, and Measurements* (Kluwer Academic/Plenum Publishers, New York, 2001).
- [8] P.W. Shor, in *Proceedings of the 35th Annual Symposium on the Foundations of Computer Science*, edited by S. Goldwasser (IEEE Computer Society Press, Los Alamos, Ca, 1994).
- [9] J.G. Rarity and P.R. Tapster, Phys. Rev. A **59**, R35 (1999).
- [10] D.M. Greenberger, M. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe* editor M. Kafatos (Kluwer, Dordrecht, 1989).
- [11] K. Wódkiewicz, Liwei Wang, and J.H. Eberly, Phys. Rev. A **47**, 3280 (1993).
- [12] C.J. Hood, W. Lange, H. Mabichi, and H.J. Kimble Phys. Rev. Lett. **10** 4710 (1995).
- [13] C. Monroe, D.M. Meekhof, B.E. King, W.M. Itano, and D.J. Wineland, Phys. Rev. Lett. **75**, 4714 (1995).
- [14] M. Weidinger, B.T.H. Varcoe, R. Heerland, and H. Walther Phys. Rev. Lett. **82** 3795 (1999); S. Haroche, AIP Conf. Proc. **464**, 45 (1999).
- [15] G. Rempe, Ann Phys. (Leipzig) **9** 843 (2000).
- [16] B. Bederson and H. Walther, editors *Advances in Atomic, Molecular, and Optical Physics*, Vol. **42** (Academic Press, New York, 2000).
- [17] B. Julsgaard, A. Kozhekin, and E. Polzik, Nature **413**, 400 (2001).
- [18] M.B. Plenio, S.F. Huelga, A. Beige, and P.L. Knight, Phys. Rev. A **59**, 2468 (1999).
- [19] A. Vourdas, Phys. Rev. A **41**, 1653 (1990).
- [20] A.S. Shumovsky and Ö.E. Müstecaplıoğlu, J. Mod. Optics, **45**, 619 (1998).
- [21] A.S. Shumovsky, J. Phys. A **32**, 6589 (1999).
- [22] A.S. Shumovsky, in *Modern Nonlinear Optics, Part I, Second Edition, Advances in Chemical Physics*, Vol. 119. Edited by M.W. Evans. Series Editors I. Prigogine and S.A. Rice (Wiley New York, 2001).
- [23] A.A. Klyachko and A.S. Shumovsky, *Entangled States and Local Measurements*, arXiv:quant-ph/0203099.
- [24] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **80**, 5239 (1998).
- [25] M.B. Plenio and V. Verdal, Contemp. Phys. **39**, 431 (1998).
- [26] D.M. Greenberger, M. Horne, A. Simony, and A. Zeilinger, Am. J. Phys. **58**, 1131 (1990).
- [27] N.D. Mermin, Phys. Rev. Lett. **65**, 1838 (1990).
- [28] A.V. Belinskii and D.N. Klyshko, Phys. Usp. **36**, 460 (1993).
- [29] R. Horodecki, P. Horodecki, and M. Horodecki M., Phys. Lett. A **200**, 340 (1995).
- [30] A. Peres, Found. Phys. **29**, 589 (1999).
- [31] R.F. Werner and M.M. Wolf, Phys. Rev. A **64**, 010102(R) (2001).
- [32] Č. Brukner, M. Žukovski, and A. Zeilinger, *The Essence of Entanglement*, arXiv:quant-ph/0106119v1.
- [33] J.F. Clauser, M.A. Horne, A. Shimoni, and R.A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
- [34] M.O. Scully and M.S. Zubairy, *Quantum Optics* (Cambridge University Press, New York, 1997).
- [35] E.g., see: N.N. Bogolubov Jr, B.I. Sadovnikov, and A.S. Shumovsky, *Mathematical Methods of Statistical Mechanics of Model Systems* (CRC Press, Boca Ration, FL, 1994).
- [36] J.I. Cirac, Nature **409**, 63 (2001).
- [37] A. Vourdas, Phys. Rev. A **65**, 042321 (2002).
- [38] A.S. Shumovsky and Ö.E. Müstecaplıoğlu, Phys. Rev. Lett. **80**, 1202 (1998).
- [39] See: A.S. Shumovsky, in Ref. [7].
- [40] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, New York, 1995).
- [41] A.V. Burlakov, M.V. Chekhova, O.A. Karabutova, D.N. Klyshko, and S.P. Kulik, Phys. Rev. A **60**, R4209 (1999).
- [42] H. Bechmann-Pasquinucci and A. Peres, Phys. Rev. Lett. **85**, 3313 (2000).
- [43] C.K. Law and J.H. Eberly, Phys. Rev. A **47**, 3195 (1993); R.R. Puri, C.K. Law, and J.H. Eberly, Phys. Rev. A **50**, 4212 (1994); C. Cabrillo, J.I. Cirac, P. Garsia-Fernandez, and P. Zoller, Phys. Rev. A **59**, 1025 (1999); M. Hennrich, T. Legero, K. Khun, and G. Rempe, Phys. Rev. Lett. **85**, 4872 (2000); A. Beige, W.J. Munro, and P.L. Knight, Phys. Rev. A **62**, 052102 (2000).

FIG. 1. Atomic Raman-type interaction with pump (P) and Stokes (S) photons.

